
The slide has a vertical yellow bar on the left with a photograph of a large, ornate metal key standing upright on a gravel surface. The rest of the slide has a blue background. The title "Airway Devices" is in yellow text. Below it is a bullet point in white text. At the bottom, there is a footer with three items: a date, a title, and a page number.

**Airway Devices**

- ♦ **This section is intended for additional clarification and pictures of devices.**

August 2002      Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force      2




## Airway Devices

- ♦ It also includes several studies relative to airway device utilization.


August 2002

Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force

3



## Oropharyngeal Airways



*Guedel plastic oropharyngeal airway.*  
Reproduced by courtesy of the *Collection of Anesthesia and Intensive Care Medicine at the Institute for the History of Medicine in Vienna (Austria).*


*Ernst Zadrobilek, MD, (Vienna, Austria),*

August 2002

Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force

4

# Nasopharyngeal Airways



Wendl red rubber nasopharyngeal airway (see Figures 1 to 4) from the [Collection of Anesthesia and Intensive Care Medicine at the Institute for the History of Medicine in Vienna \(Austria\)](#). Ernst Zadrobilek, MD, (Vienna, Austria)

August 2002

Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force

5


# Mounted Suction




August 2002

Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force

6



## Portable Suction



August 2002

Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force

7



## ECT Kit




August 2002

Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force

8

# ECT Procedure Outline

1. Cross section:



The diagram shows a cross-section of an ECT device. It consists of two main channels, labeled 1 and 2, which are curved around a yellow, oval-shaped object. Channel 1 is the outer curve, and channel 2 is the inner curve. The device is shown in a cross-section view, revealing the internal structure of the channels.


August 2002

Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force

9

# ECT Procedure Outline

2. Insertion:




The diagram illustrates the insertion of an ECT device into a patient's airway. A hand is shown holding the device, which is being inserted into the patient's mouth. The device is shown in a cross-section view, revealing the internal structure of the channels. The patient's head is shown in profile, with the airway highlighted in red. The device is shown being inserted into the airway, with the hand positioned to guide it.

August 2002


Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force

10



# ECT Procedure Outline


3. Inflation of balloon and cuff:



August 2002


Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force

11



# ECT Procedure Outline

4. Esophageal placement and ventilation:



August 2002

Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force

12

## ECT Procedure Outline

### 5. Tracheal placement and ventilation:

August 2002

Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force

13


## ECT Advantages

- ♦ **Non invasive**
  - Helpful under difficult circumstances with respect to space and illumination
  - No preparation necessary
  - Blind insertion possible, however use laryngoscope whenever feasible !
  - Simultaneous fixation after inflation of oropharyngeal balloon
  - Works in tracheal or esophageal position
  - Minimized risk of aspiration
  - Application of high ventilatory pressures possible
  - Independent of power supply
  - Optimal method in emergency intubation and in cases of bleeding when visualization of vocal cords is impossible

August 2002

Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force


14



## ECT Contraindications

- Patients with intact gag reflexes
- Patient's height below 4 feet
- Patients with known esophageal pathology
- Patients after ingestion of caustic substances
- Central-airway obstruction


August 2002 Kansas Airway Supplement Kansas BEMS EMS Educator Task Force 15



## Airway Related Research Selected Studies

August 2002 Kansas Airway Supplement Kansas BEMS EMS Educator Task Force 16






**Ventilations Must be...**

**“adequately monitored, including continuous monitoring of end-tidal carbon dioxide concentrations.”**


Prehosp Emerg Care 2001  
Jan-Mar;5(1):73-8

August 2002 Kansas Airway Supplement Kansas BEMS EMS Educator Task Force 17



**Data were recorded by the EMS provider on 167 [corrected] adult patients (age range 16-92 years) in whom a PTL or endotracheal (ET) airway insertion was attempted.**

August 2002 Kansas Airway Supplement Kansas BEMS EMS Educator Task Force 18




**Data were recorded by the EMS provider on 167 [corrected] adult patients (age range 16-92 years) in whom a PTL or endotracheal (ET) airway insertion was attempted.**

**RESULTS:** There were no significant differences between the PTL and ET groups, either in patient demographics or in rates of successful ventilation with either airway overall or in trauma-related versus non-trauma-related cases, male versus female patients, or volunteer versus paid EMS providers.

Prehospital Disaster Med 1992 Jan-Mar;7(1):13-8  
McMahan S, Ornato JP, Racht EM, Cameron J.


August 2002 Kansas Airway Supplement Kansas BEMS EMS Educator Task Force 19



**1139 patients were resuscitated with the ETC and the semiautomated external defibrillator as part of the CPR protocol for prehospital management of cardiac arrest by basic emergency medical technicians.**

Can J Anaesth 1998 Jan;45(1):76-80  
Vezina D, Lessard MR, Bussieres J, Topping C, Trepanier CA.

August 2002 Kansas Airway Supplement Kansas BEMS EMS Educator Task Force 20



**1139 patients were resuscitated with the ETC and the semiautomated external defibrillator as part of the CPR protocol for prehospital management of cardiac arrest by basic emergency medical technicians.**


**Eight of these patients presented with subcutaneous emphysema.**

***In addition:***

**These cases suggest that subcutaneous emphysema, pneumomediastinum and pneumoperitoneum might be complications associated with the use of the ETC.**

Can J Anaesth 1998 Jan;45(1):76-80  
Vezina D, Lessard MR, Bussieres J, Topping C, Trepanier CA.

August 2002 Kansas Airway Supplement Kansas BEMS EMS Educator Task Force 21




**End-tidal CO<sub>2</sub> (ETCO<sub>2</sub>) Detector (EASY CAP)**

**This detector was used for 121 patients during CPR with a laryngeal mask airway or face mask by authorized emergency lifesaving technicians.**

Am J Emerg Med 1999 Mar;17(2):203-6  
Nakatani K, Yukioka H, Fujimori M, Maeda C, Noguchi H, Ishihara S, Yamanaka I, Tase C.

August 2002 Kansas Airway Supplement Kansas BEMS EMS Educator Task Force 22




## End-tidal CO<sub>2</sub> (ETCO<sub>2</sub>) Detector (EASY CAP)

**At 7 to 15 minutes after the  
initiation of CPR, ETCO was:**

<b>&lt;0.5% in 30 cases</b>	<b>(group A)</b>
<b>0.5% to 2.0% in 46 cases</b>	<b>(group B)</b>
<b>&gt;2.0% in 45 cases</b>	<b>(group C)</b>

August 2002      Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force

23



## End-tidal CO<sub>2</sub> (ETCO<sub>2</sub>) Detector (EASY CAP)


**At 7 to 15 minutes after the initiation of  
CPR, ETCO was:**

**The rate of return of  
spontaneous circulation  
was:**

<b>&lt;0.5% in 30 cases</b>	<b>(group A) 17%</b>
<b>0.5% to 2.0% in 46 cases</b>	<b>(group B) 24%</b>
<b>&gt;2.0% in 45 cases</b>	<b>(group C) 48%</b>

August 2002      Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force


24



The most appropriate airway device for use in EMS systems staffed by basic skilled EMTs with (EMT-Ds) or without (EMT-Bs) defibrillation capabilities is *still a matter of debate.*

Resuscitation 2002 Jan;52(1):77-83  
Lefrancois DP, Dufour DG.


August 2002 Kansas Airway Supplement Kansas BEMS EMS Educator Task Force 25



The most appropriate airway device for use in EMS systems staffed by basic skilled EMTs with (EMT-Ds) or without (EMT-Bs) defibrillation capabilities is \_\_\_\_\_.

Resuscitation 2002  
Jan;52(1):77-83  
Lefrancois DP, Dufour DG


August 2002 Kansas Airway Supplement Kansas BEMS EMS Educator Task Force 26



**The purpose of this study was to assess the feasibility, safety and effectiveness of the Esophageal Tracheal Combitube (ETC) when used by EMT-Ds in cardiorespiratory arrest patients of all etiologies.**

Resuscitation 2002  
Jan;52(1):77-83  
Lefrancois DP, Dufour DG.

August 2002 Kansas Airway Supplement Kansas BEMS EMS Educator Task Force 27




**831 Arrest Victims Studied  
Esophageal Tracheal Combitube (ETC)**

**Placement attempted in 760 patients**

Resuscitation 2002  
Jan;52(1):77-83  
Lefrancois DP, Dufour DG.

August 2002 Kansas Airway Supplement Kansas BEMS EMS Educator Task Force 28




**831 Arrest Victims Studied  
Esophageal Tracheal Combitube (ETC)**

**Placement attempted in 760 patients  
Placement was successful in 725  
(95.4%)**

Resuscitation 2002  
Jan;52(1):77-83  
Lefrancois DP, Dufour DG.

August 2002 Kansas Airway Supplement Kansas BEMS EMS Educator Task Force 29




**831 Arrest Victims Studied  
Esophageal Tracheal Combitube (ETC)**

**Placement attempted in 760 patients  
Placement was successful in 725  
(95.4%)  
Ventilation was successful in 695  
(91.4%)**

Resuscitation 2002  
Jan;52(1):77-83  
Lefrancois DP, Dufour DG.


August 2002 Kansas Airway Supplement Kansas BEMS EMS Educator Task Force 30



**It is essential that airway devices be constantly and carefully monitored for placement throughout their use.**

Prehosp Emerg Care 1999  
Oct-Dec;3(4):273-8  
Falk JL, Sayre MR

August 2002 Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force 31




**A prospective, controlled study to evaluate the difficulty and complications of insertion, recognition of esophageal versus tracheal placement, skill proficiency, and retention.**

Ann Emerg Med 1993 Aug;22(8):1263-8  
Atherton GL, Johnson JC.

August 2002 Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force 32





**Fifty-two cases of paramedic prehospital  
Combitube insertion were examined.**

Ann Emerg Med 1993 Aug;22(8):1263-8  
Atherton GL, Johnson JC.


August 2002 Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force 33



**Combitube insertion was attempted on 52  
prehospital patients in cardiac arrest**

Ann Emerg Med 1993 Aug;22(8):1263-8  
Atherton GL, Johnson JC.

August 2002 Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force 34



**Paramedics recognized  
esophageal versus tracheal  
placement in 100% of the cases.**

Ann Emerg Med 1993 Aug;22(8):1263-8  
Atherton GL, Johnson JC.


August 2002 Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force 35



**The Combitube was inserted  
successfully into 64% of the  
patients who could not be  
endotracheally intubated by the  
conventional visualized method.**

Ann Emerg Med 1993 Aug;22(8):1263-8  
Atherton GL, Johnson JC.

August 2002 Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force 36



**The Combitube was inserted successfully 71% of the time when used as a first-line airway adjunct.**

Ann Emerg Med 1993 Aug;22(8):1263-8  
Atherton GL, Johnson JC.


August 2002 Kansas Airway Supplement Kansas BEMS EMS Educator Task Force 37



**A follow-up study on 11 randomly selected paramedics involved in the field study was conducted 15 months later.**

Ann Emerg Med 1993 Aug;22(8):1263-8  
Atherton GL, Johnson JC.


August 2002 Kansas Airway Supplement Kansas BEMS EMS Educator Task Force 38



**Nine of 11 the paramedics  
demonstrated inadequate skill  
retention in the follow-up study.**

Ann Emerg Med 1993 Aug;22(8):1263-8  
Atherton GL, Johnson JC.

August 2002 Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force 39



**Nine of 11 the paramedics  
demonstrated inadequate skill  
retention in the follow-up study.**

Ann Emerg Med 1993 Aug;22(8):1263-8  
Atherton GL, Johnson JC.

August 2002 Kansas Airway Supplement  
Kansas BEMS EMS Educator Task Force 40